Haemonchus contortus in the UK

Results of the Survey


January 2024

Lesley Stubbings and Nerys Wright



What is your main role?








Geographical representation












How would you describe the change in incidence of clinical disease


- by occupation



Ewes




Lambs



In your opinion, how serious is the impact of haemonchus on sheep farms? (1 = Not at all serious; 10 = Extremely serious)






How serious is the impact of haemonchus on sheep farms, by occupation









Do you think there has been a shift in the time of year when you see problems with haemonchus?






Do you think there has been a shift in the time of year, by occupation





If you answered 'Yes' above, can you say what the change has been?






What is the 'change'?




65 provided a rationale to their response Other:

- Finding late summer to October a problem
- More sporadic and more variable from spring in ewes (overwintered L4?) to late autumn in adults and lambs
- Earlier in the year on farms that had a problem the previous year
- New regions
- Earlier and later in the year and for longer (x2)
- Weather related personnel experience more active when hot
- Winter




What diagnostic/monitoring tool do you use for haemonchus (tick all that apply)






In your experience, what level of resistance have you found to the different anthelmintic groups in haemonchus?






What level of resistance have you found – by opinion






Have you used or know of clients who have used Barbervax?






There were several questions that were categorised into these headings.

Questions from the respondents







Best strategy/best
wormer/most
effective/first ones to
use/when to use a
wormer/rotating
wormers/

Can we treat arrested larvae in sheep successfully with a winter closantel dose?

Whole flock treatment
Vs selective (ie
symptoms/thin/
FAMACHA)

The issue of fluke on the farm also – wrong diagnosis/use of closantel/

Is it best to use moxidectin for longer cover

When to treat/move most effectively as not spread the worm to new pastures

Are there any other chemical groups that we could bring in and licence from abroad?

Is there a likely new treatment on the horizon

Further information

- This survey was funded by SCOPS <u>www.scops.org.uk</u>
- If you wish to utilise this information, please reference 'SCOPS Haemonchus survey, 2024'
- Please contact <u>enquiries@nationalsheep.org.uk</u> for further information on this topic. The National Sheep Association provides secretarial support for SCOPS and will direct your enquiry appropriately.

